

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Northern Blot: 7 Bacterial Species Probed with Conserved 22mer from 4.5S



Hybridized with Ad4.5S13Vnf:

- RT
- 120V
- 5% gel
- 8M urea

Fig. 1

Northern Blot to compare abundance of 5S and 4.5S





Fig. 2

Detection of 4.5S RNA in Cultured E.coli by Adapter-Capture and Fluorescent Sandwich Probe



3λ /20 RNA from 1 ml of overnight culture

Fig. 3

In-Gel Strand Displacement




Fig. 4

Adapters for analysis of RNA on Universal Capture Gels.




Fig. 5




Fig. 6




Fig. 7